

AD-A048 152

CLEMSON UNIV S C DEPT OF MATHEMATICAL SCIENCES  
ON A GENERALIZATION OF FREUND'S BIVARIATE FAILURE MODEL.(U)  
JUN 77 T J TOSCH, P T HOLMES

F/G 12/1

N00014-75-C-0451

UNCLASSIFIED


N86

NL

| OF |  
AD  
A048152



END  
DATE  
FILMED  
1-78  
DDC



NATIONAL BUREAU OF STANDARDS  
MICROCOPY RESOLUTION TEST CHART

AD A048152



10  
B.S.

DEPARTMENT  
OF  
MATHEMATICAL  
SCIENCES

CLEMSON UNIVERSITY  
Clemson, South Carolina



AD No. \_\_\_\_\_  
DDC FILE COPY



6  
ON A GENERALIZATION OF FREUND'S  
BIVARIATE FAILURE MODEL

BY

10 THOMAS J. TOSCH\*  
PAUL T. HOLMES

9 Technical rept.

DEPARTMENT OF MATHEMATICAL SCIENCES  
CLEMSON UNIVERSITY

11 1 JUNE 1977


14 N86, TR-252  
N86



15  
RESEARCH SUPPORTED BY THE OFFICE OF NAVAL RESEARCH  
TASK NR 042-271 CONTRACT N00014-75-C-0451

REPRODUCTION IN WHOLE OR PART IS PERMITTED FOR ANY PURPOSE OF  
THE U. S. GOVERNMENT, DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.

\*NOW AT THE UNIVERSITY OF WASHINGTON



407 183 mt

### Abstract

A bivariate failure model is proposed in which the residual lifetime of one component is dependent on the working status of the other. General properties of the model are discussed, and the maximum likelihood estimates of the parameters are found in a bivariate exponential-like special case.

Keywords: Bivariate failure model, bivariate exponential distribution, maximum likelihood estimation.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION _____	
BY _____	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	



## 1. INTRODUCTION

A new bivariate failure model is proposed in which the residual lifetime of one component is dependent on the working status of the other component. This is applicable when the failure of one component puts more (possibly less) strain on the remaining components, for example, the kidneys. Section two derives properties of the lifetimes, including their joint Laplace-Stieltjes transform.

In the third section a bivariate exponential-like special case is considered. In this example maximum likelihood estimates of the parameters are obtained and their asymptotic distribution studied. This model is compared with the bivariate exponential models of Freund [3] and Marshall and Olkin [4].

## 2. MODEL DEFINITION AND GENERAL PROPERTIES

Label the two components of the system A and B with lifetimes S and T respectively. The lifetimes of the two components are dependent, in that the failure of one component affects the residual lifetime of the other. To describe S and T, let X, Y, U, V be non-negative mutually independent random variables with X and Y absolutely continuous. Then we write

$$\begin{aligned} S &= \min(X, Y) + U \cdot I_{\{X > Y\}} \\ T &= \min(X, Y) + V \cdot I_{\{X \leq Y\}} \end{aligned} \tag{2.1}$$

We will obtain an expression for the joint survival distribution

$\bar{F}(s,t) = \Pr\{S > s, T > t\}$ . Assuming  $s < t$  and conditioning on the values of  $X$  and  $Y$ , we obtain

$$\bar{F}(s,t) = \int_0^\infty \int_0^\infty \Pr\{S > s, T > t | X=x, Y=y\} dF_X(x) dF_Y(y).$$

Now partition the region  $[0,\infty) \times [0,\infty)$  into 12 subregions based on the relative sizes of  $x,y,s,t$ . The only non-zero contributions are  $\Pr\{V, t-x\}$  on the region  $[s,t] \times [x,\infty)$  and 1 on the region  $[t,\infty) \times [t,\infty)$ . Thus the following can be established.

Theorem 1:

$$\bar{F}(s,t) = \begin{cases} \bar{F}_X(t)\bar{F}_Y(t) + \int_s^t \bar{F}_V(t-x)\bar{F}_Y(x) dF_X(x) & \text{if } s < t \\ \bar{F}_X(s)\bar{F}_Y(s) & \text{if } s = t \\ \bar{F}_X(s)\bar{F}_Y(s) + \int_t^s \bar{F}_U(s-y)\bar{F}_X(y) dF_Y(y) & \text{if } s > t. \end{cases}$$

The joint Laplace-Stieltjes transform of  $(S,T)$  is defined to be  $f^*(a,b) = \int_0^\infty \int_0^\infty e^{-as-bt} F(ds,dt)$ , where  $F(ds,dt)$  is the measure determined by the survival function  $\bar{F}(s,t)$ . This measure can be represented by

$$F(ds,dt) = \begin{cases} \bar{F}_Y(s) dF_V(t-s) dF_X(s) & \text{if } s < t \\ \bar{F}_Y(s) p_V dF_X(s) + F_X(s) p_U dF_Y(s) & \text{if } s = t \\ \bar{F}_X(t) dF_U(s-t) dF_Y(t) & \text{if } s > t, \end{cases} \quad (2.2)$$

where  $p_V = \Pr\{V=0\}$  and  $p_U = \Pr\{U=0\}$ . This expression is used to evaluate  $f^*(a,b)$  and we get

Theorem 2:

$$f^*(a, b) = f_V^*(b) \int_0^\infty e^{-(a+b)s} \bar{F}_Y(s) dF_X(s) + f_U^*(a) \int_0^\infty e^{-(a+b)s} \bar{F}_X(s) dF_Y(s).$$

These integrals cannot be evaluated in general, but can be for certain important special cases. In particular we have

Corollary 3: If Y has an exponential distribution with parameter  $\beta$ , then

$$f^*(a, b) = f_V^*(b) f_X^*(a+b+\beta) + (f_U^*(a) \beta / (a+b+\beta)) [1 - f_X^*(a+b+\beta)].$$

Moments can be calculated by differentiation using this expression.

### 3. ESTIMATION IN A SPECIAL CASE

As a special case suppose that X and Y are exponentially distributed with parameters  $\alpha$  and  $\beta$  respectively,  $\Pr\{U > t\} = qe^{-\alpha't}$ ,  $\Pr\{V > t\} = qe^{-\beta't}$ , where  $\alpha, \beta, \alpha', \beta' > 0$ ,  $0 \leq q \leq 1$ ,  $t \geq 0$ . Freund's [3] model corresponds to the case  $q=1$ . The parameter  $q$  allows for simultaneous failure of the components, since  $\Pr\{S=T\} = 1-q = p$ . Using results from section 2, properties (including moments) can be derived. In particular,

$$\bar{F}(s, t) = \begin{cases} e^{-(\alpha+\beta)t} + \frac{q\alpha e^{-\beta't}}{\alpha+\beta-\beta'} [e^{-(\alpha+\beta-\beta')s} - e^{-(\alpha+\beta-\beta')t}] & \text{if } s < t \\ e^{-(\alpha+\beta)s} & \text{if } s = t \\ e^{-(\alpha+\beta)s} + \frac{q\beta e^{-\alpha's}}{\alpha+\beta-\alpha} [e^{-(\alpha+\beta-\alpha')t} - e^{-(\alpha+\beta-\alpha')s}] & \text{if } s > t. \end{cases}$$

(3.1)



Marshall and Olkin [4] defined a bivariate lack of memory property by  $\Pr\{S > s+\Delta, T > t+\Delta\} = \Pr\{S > s, T > t\} \cdot \Pr\{S > \Delta, T > \Delta\}$  for all  $s, t, \Delta \geq 0$ . The survival distribution (3.1) possesses this property and has mixtures of exponential distributions as marginals.

The measure determined by (3.1) however is not absolutely continuous. If we let  $\mu_i$  represent Lebesgue measure on  $R_i$ , then the measure defined by  $\mu(A) = \mu_2(A) + \mu_1\{x: (x, x) \in A\}$  is a suitable dominating measure for maximum likelihood estimation. This is the same dominating measure used by Bhattacharyya and Johnson [1].

Consider a sample of  $N$  independent observations on  $(S, T)$ ,  $\{(S_1, t_1), \dots, (S_N, t_N)\}$ . The following notation simplifies the likelihood function. Let  $A_1 = \{(S_i, t_i) | s_i < t_i\}$ ,  $A_2 = \{(S_i, t_i) | s_i > t_i\}$  and  $A_3 = \{(S_i, t_i) | s_i = t_i\}$ . Further let  $N_1 = \#A_1$ ,  $N_2 = \#A_2$ ,  $N_3 = \#A_3$ ,  $S_1 = \sum_{A_1} s_i$ ,  $S_2 = \sum_{A_2} s_i$ ,  $R = \sum_{A_3} s_i$ ,  $T_1 = \sum_{A_1} t_i$ ,  $T_2 = \sum_{A_2} t_i$ . Here  $\#A$  is the number of items in  $A$ . Then  $L$ , the likelihood function, can be expressed as

$$L = (\alpha\beta')^{N_1} (1-p)^{N_1+N_2} p^{N_3} (\alpha+\beta)^{N_3} (\beta\alpha')^{N_2} \exp[-(\alpha+\beta)(S_1+R+T_2) - \alpha'(S_2-T_2) - \beta'(T_1-S_1)]. \quad (3.2)$$

Solving for the maximum likelihood estimates, one obtains

Theorem 4:

- (i) If  $N_1=N_2=0$ , then  $\hat{p}=1$  and  $\alpha, \beta, \alpha', \beta'$  cannot be estimated,

- (ii) If  $N_1=0$ , but  $N_2 \neq 0$ , then  $\hat{p}=N_3/N$ ,  $\hat{\alpha}=0$ ,  $\hat{\beta}=N/(S_1+R+T_2)$ ,  $\hat{\alpha}'=N_2/(S_2-T_2)$ , and  $\hat{\beta}'$  cannot be estimated.
- (iii) If  $N_1 \neq 0$ , but  $N_2=0$  then  $\hat{p}=N_3/N$ ,  $\hat{\beta}=0$ ,  $\hat{\alpha}=N/(S_1+R+T_2)$ ,  $\hat{\beta}'=N_1/(T_1-S_1)$ , and  $\hat{\alpha}'$  cannot be estimated.
- (iv) If  $N_1 \neq 0$  and  $N \neq 0$  then
- $$\hat{\alpha} = (N/S_1 + R + T_2) (N_1/N_1 + N_2)$$
- $$\hat{\beta} = (N/S_1 + R + T_2) (N_2/N_1 + N_2)$$
- $$\hat{\alpha}' = N_2/(S_2 - T_2)$$
- $$\hat{\beta}' = N_1/(T_1 - S_1)$$
- $$\hat{p} = N_3/N.$$

It is easy to obtain the biases in these estimators. Using a Lehmann, Scheffé partitioning operation (cf. Zacks [5], p. 50) it can be shown that

Theorem 5: The vector  $(N_1, N_2, S_1+R+T_2, S_2-T_2, T_1-S_1)$ , (and thus the vector of maximum likelihood estimators) is a minimal sufficient statistic of the sample  $\{(s_1, t_1), \dots, (s_N, t_N)\}$ .

To obtain the asymptotic distribution of the maximum likelihood estimates we need to restrict the parameter space as follows. Let  $\Omega^* = \{(\alpha, \beta, \alpha', \beta', p) \mid 0 < t_1 < \alpha < M_1, \dots, 0 < t_5 < p < M_5 < 1\}$ . On this space the regularity conditions presented by Chanda [2, p. 56] are satisfied. Thus we have

Theorem 6:

- (i)  $(\hat{\alpha}, \hat{\beta}, \hat{\alpha}', \hat{\beta}', \hat{p}) \rightarrow (\alpha, \beta, \alpha', \beta', p)$  as  $N \rightarrow \infty$  with probability one.

(ii)  $N^{1/2}(\hat{\alpha}-\alpha, \hat{\beta}-\beta, \hat{\alpha}'-\alpha', \hat{\beta}'-\beta', \hat{p}-p)$  is asymptotically distributed as multivariate normal with mean  $\underline{0}$  and covariance matrix  $\Sigma$ , where

$$\Sigma^{-1} = \begin{bmatrix} \frac{\alpha+\beta-p\beta}{\alpha(\alpha+\beta)^2} & \frac{p}{(\alpha+\beta)^2} & 0 & 0 & 0 \\ \frac{p}{(\alpha+\beta)^2} & \frac{\alpha+\beta-p\alpha}{\beta(\alpha+\beta)^2} & 0 & 0 & 0 \\ 0 & 0 & \frac{(1-p)\beta}{(\alpha')^2(\alpha+\beta)} & 0 & 0 \\ 0 & 0 & 0 & \frac{(1-p)\alpha}{(\beta')^2(\alpha+\beta)} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{p(1-p)} \end{bmatrix}$$

The parameters can also be estimated by maximum likelihood for a censored sample.

We see that this model generalizes Freund's model to include simultaneous failure of both components. It differs from the model of Marshall and Olkin in that the residual lifetime of one component is not independent of the status of the other component. We feel that these features will aid in the application of the model.

## References

1. Bhattacharyya, G. K. and Johnson, R. A., "Maximum Likelihood Estimation and Hypothesis Testing in the Bivariate Exponential Model of Marshall and Olkin," Technical Report No. 276, Department of Statistics, University of Wisconsin, August, 1971.
2. Chanda, K. C., "A Note on the Consistency and Maxima of the Roots of Likelihood Equations," Biometrika, 49(1954), 56-61.
3. Freund, John E., "A Bivariate Extension for the Exponential Distribution," Journal of the American Statistical Association, 56(Dec. 1961), 971-977.
4. Marshall, Albert W., and Olkin, Ingram, "A Multivariate Exponential Distribution," Journal of the American Statistical Association, 62(March 1967), 30-44.
5. Zacks, S., The Theory of Statistical Inference, New York: John Wiley and Sons, Inc., 1971.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER N86 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  On a Generalization of Freund's Bivariate Failure Model		5. TYPE OF REPORT & PERIOD COVERED  Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  Thomas J. Tosch Paul T. Holmes		8. CONTRACT OR GRANT NUMBER(s)  N00014-75-C-0451 ✓
9. PERFORMING ORGANIZATION NAME AND ADDRESS  Clemson University Dept. of Mathematical Sciences ✓ Clemson, South Carolina 29631		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  NR 042-271
11. CONTROLLING OFFICE NAME AND ADDRESS  Office of Naval Research Code 436 Arlington, Va. 22217		12. REPORT DATE June 1, 1977
		13. NUMBER OF PAGES 6
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Bivariate failure model, bivariate exponential distribution, maximum likelihood estimation.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  A bivariate failure model is proposed in which the residual lifetime of one component is dependent on the working status of the other. General properties of the model are discussed, and maximum likelihood estimates of the parameters are found in a bivariate exponential-like case.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)